



Areas bounded by the curves

Vikasana - CET 2012



In this chapter, while calculating the definite integral as the '*limit of the sum*'. We have learnt the process of finding the area bounded by the curve $y=f(x)$, the x -axis and the ordinates $x=a$ and $x=b$.




In this chapter we shall discuss the use of definite integrals. In computing areas bounded by simple curves such as straight lines, circles, parabolas and other conics.

Vikasana - CET 2012



Let $y=f(x)$ be a finite and continuous curve in the interval $[a,b]$. Then the area between the curve $y=f(x)$, x -axis and two ordinates at the points $x = a$ and $x = b$ is given by,

$$A = \int_a^b y dx = \int_a^b f(x) dx$$




Vikasana - CET 2012



Let $y=f(x)$ be a continuous curve below the x -axis. Then the area between the curve $y=f(x)$, x -axis and the ordinates $x=a$ and $x=b$ is given by

$$A = \int_a^b -y dx = - \int_a^b f(x) dx$$


$$A = \left| \int_a^b f(x) dx \right|$$





The area bounded by the curve $x=f(y)$, y -axis and the lines $y=c$ and $y=d$ ($c < d$) is given by

$$A = \int_c^d x dy = \int_c^d f(y) dy$$






If the curve $x=f(y)$ lies to the left of y -axis then the area bounded by the curve $y=f(x)$ and the lines $y=c$ and $y=d$ is given by

$$A = \int_c^d (-x) dy = - \int_c^d x dy$$

$$A = \left| \int_c^d f(y) dy \right|$$




Vikasana - CET 2012



If the curve crosses x -axis at one point 'C' then the area bounded by the curve is given by.


$$A = \left| \int_a^c f(x) dx \right| + \left| \int_c^b f(x) dx \right|$$





If the curve crosses x -axis in two points **c & d** , then the area between the curve $y=f(x)$, the **x -axis** and the ordinates $x=a$ & $x=b$ is


$$A = \left| \int_a^c f(x) dx \right| + \left| \int_c^d f(x) dx \right| + \left| \int_d^b f(x) dx \right|$$





The area enclosed between the curves $y=f_1(x)$ and $y=f_2(x)$ between the ordinates $x=a$ & $x=b$ is given by

$$\int_a^b |f_1(x) - f_2(x)| dx$$





If the two curves do not cross each other between lines $x=a$ & $x=b$, then the area is

$$\left| \int_a^b f_1(x) dx - \int_a^b f_2(x) dx \right|$$



Curve Sketching for Area

For the evaluation of area of bounded regions, it is very essential to draw the rough sketch of the curves. The following points are very useful to draw a rough sketch of the curve.



- For all ' x ' for which $y=f(x)=0$ ($a \leq x \leq b$)
- Mark these points on x - *axis*.
- In case of two curves, find the point of intersection of two curves.
- Use symmetry of the curve in finding area.

Vikasana - CET 2012



Symmetry about x -axis –

If the equation of the curve does not change when ' y ' is changed to ' $-y$ ', then the curve is symmetrical about x - axis.

(i.e. If only even power of ' y ' occur, then the curve is symmetrical about x -axis).

Ex: $y^2 = 4ax$ is symmetrical about x -axis.

Vikasana - CET 2012



Symmetry about y -axis :

If the equation of the curve does not change, when x is changed to $-x$, then the curve is symmetrical about y -axis. (If only even power of x occur in the equation then the curve is symmetrical about y -axis)

Ex: $x^2=4ay$ is symmetrical about y -axis.



Symmetry in opposite quadrants :

If on replacing x by $-x$ and y by $-y$. Then the equation of the curve does not change (remains same). Then the curve is symmetric in opposite quadrants.

Ex: $y = \sin x$ is symmetrical in opposite quadrants.



Symmetric about the line $y = x$:

If the equation of the curve remains same on interchanging x and y , then the curve is symmetrical about the line $y=x$.

Ex: $x^3+y^3=3axy$ is symmetrical about the line $y=x$.



Some standard results on area :

- The area of the region bounded by $y^2=4ax$ and $x^2=4by$ is $\frac{16ab}{3}$ sq units.
- Area of the region bounded by $y^2=4ax$ and $y=mx$ is $\frac{8a^2}{3m^3}$ sq units.
- Area of the region bounded by $y^2=4ax$ and its latus rectum is $\frac{8a^2}{3}$ sq units.



- Area bounded by $y = \sin x$, x -axis is 2 sq units. Infact, area of one loop of $y = \sin x$ and $y = \cos x$ is 2sq. units
- Area bounded by, $y = \log_e x$, $y = 0$ and $x = 0$ is 1sq units



- Area of region bounded by the curve $y = \sin ax$ and x -axis in $[0, n\pi]$ is $\frac{2n}{a}$
 - Area of region bounded by the curve $y = \cos ax$ and x -axis in $[0, n\pi]$ is $\frac{2n}{a}$
 - Area of region bounded by one arch of $\sin ax$ or $\cos ax$ and x -axis is $\frac{2}{a}$ sq units.
 - Area of circle $x^2 + y^2 = a^2$ is πa^2 sq. units
- Vikasana - CET 2012



- The area of region bounded by parabola $y=ax^2+bx+c$ or $x=ay^2+by+c$ & x -axis is $\frac{(b^2-4ac)^{\frac{3}{2}}}{6a^2}$
- The area ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is πab sq units.



1. The area region bounded by the parabolas $y^2=4ax$ and $x^2=4ay$ is

a) $\frac{16a^2}{3}$

b) $\frac{32a^2}{3}$

c) $\frac{9a^2}{2}$

d) *none*



2. The area enclosed between the parabolas $y^2=4x$ and $x^2=4y$ is

a) $\frac{3}{4}$ sq units

b) 16 sq units

c) $\frac{16}{3}$ sq units

d) $\frac{32a^2}{3}$ sq units



3. The area enclosed between the parabolas $y^2=6x$ and $x^2=6y$ is

a) 12 sq. units

b) $\frac{16}{3}$ sq. units

c) 36 sq. units

d) none of these



4. The area inside the parabola $y^2=4ax$ between the lines $x=a$ and $x=4a$ is

a) $4a^2$

b) $28a^2$

c) $\frac{28a^2}{3}$

d) $\frac{56a^2}{3}$



Since $y^2=4ax$ and is symmetrical about x -axis

Area of the region = 2(area of the region in the 1st quadrant)

$$= 2 \int_a^{4a} y dx = 2 \int_a^{4a} \sqrt{4ax} dx = 2 \cdot 2\sqrt{a} \int_a^{4a} \sqrt{x} dx$$

$$= 4\sqrt{a} \left[\frac{x^{3/2}}{3/2} \right]_a^{4a} = \frac{8}{3} \sqrt{a} \left[(4a)^{3/2} - a^{3/2} \right] = \frac{8}{3} \sqrt{a} \left[8a^{3/2} - a^{3/2} \right]$$



5. The area bounded by the parabola $y^2=4ax$ and the line $x=a$ and $x=4a$ and x -axis is

a) $\frac{35a^2}{3}$

b) $\frac{4a^2}{3}$

c) $\frac{7a^2}{3}$

d) $\frac{28a^2}{3}$



6. The area of the figure bounded by

$y = \cos x$ and $y = \sin x$ and the

ordinates $x = 0$ and $x = \frac{\pi}{4}$ is

a) $\frac{1}{2}(\sqrt{2} - 1)$

b) $\frac{1}{\sqrt{2}}$

c) $\sqrt{2} - 1$

d) $\sqrt{2} + 1$



$$\begin{aligned} \text{Required Area} &= \int_0^{\frac{\pi}{4}} (\cos x - \sin x) dx \\ &= \left[\sin x + \cos x \right]_0^{\frac{\pi}{4}} \\ &= \left(\sin \frac{\pi}{4} + \cos \frac{\pi}{4} \right) - (\sin 0 + \cos 0) \\ &= \left(\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} \right) - (0 + 1) = \frac{2}{\sqrt{2}} - 1 = \sqrt{2} - 1 \end{aligned}$$

Vikasana - CET 2012



7. The area bounded by $y = \log_e x$, the x -axis and the line $x = e$ is

a) 1

b) $1 - \frac{1}{e}$

c) $1 + \frac{1}{e}$

d) e



$$\text{Area} \int_1^e \log x dx$$

$$= [x \log x - x]_1^e$$

$$= (e \log e - e) - (1 \log 1 - 1)$$

$$= (e - e) - (0 - 1) = 1$$

$$y = \log x$$

at $x = e$

$$y = \log_e e = 1$$



8. The area of the region bounded by the parabola $y=x^2+1$ and the straight line $x+y=3$ is given by,

a) $\frac{45}{7}$

b) $\frac{25}{4}$

c) $\frac{\pi}{18}$

d) $\frac{9}{2}$




Given $y = x^2 + 1$ & $x + y = 3 \Rightarrow y = 3 - x$

ie. $3 - x = x^2 + 1$

$(x + 2)(x - 1) = 0 \Rightarrow x = 1, -2$

Required area $= \int_{-2}^1 (3 - x) - (x^2 + 1) dx$

$= \int_{-2}^1 (2 - x - x^2) dx = 2x - \frac{x^2}{2} - \frac{x^3}{3} \Big|_{-2}^1$





$$= \left(2 - \frac{1}{2} - \frac{1}{3} \right) - \left(-4 - \frac{4^2}{2} + \frac{8}{3} \right)$$

$$= \left(\frac{12 - 3 - 2}{6} \right) - \left(\frac{-18 + 8}{3} \right)$$

$$\frac{7-2-1}{6} - \left(-4 - \frac{4^2}{2} + \frac{8}{3} \right)$$

$$\frac{7-2-1}{6} - \left(-4 - \frac{4^2}{2} + \frac{8}{3} \right) = \frac{9}{2}$$

Vikasana - CET 2012



9. The area of portion of the circle $x^2+y^2=64$ which is exterior to the parabola $y^2=12x$

a) $\frac{16}{3}(4\pi - \sqrt{3})$ sq units b) $\frac{16}{3}(8\pi - \sqrt{3})$ sq units

c) $\frac{16}{3}(8 + \sqrt{3})$ sq units d) None of these



In the first quadrant the point of intersection of the circle $x^2 + y^2 = 64$ and the parabola $y^2 = 12x$ is $(4, \pm 4\sqrt{3})$

$$x^2 + y^2 = 64 \quad \text{ie., } x^2 + 12x - 64 = 0$$

$$\Rightarrow x^2 + 16x - 4x - 64 = 0 \Rightarrow (x - 4)(x + 16) = 0$$

$$\text{ie., } y^2 = 48 \quad \therefore y = \pm 4\sqrt{3}$$



Required area =

Area of the circle – Area of circle exterior to the parabola.

$$= 64\pi - 2 \int_0^4 y dx - 2 \int_4^8 y dx$$

$$= 64\pi - 2 \int_0^4 2\sqrt{3}\sqrt{x} dx - 2 \int_4^8 \sqrt{64 - x^2} dx$$


Vikasana - CET 2012



$$= 64\pi - 4\sqrt{3} \left[\frac{x^{\frac{3}{2}}}{\frac{3}{2}} \right]_0^4 - 2 \left[\frac{x}{2} \sqrt{64 - x^2} + \frac{64}{2} \sin^{-1} \left(\frac{x}{8} \right) \right]_0^4$$

$$= 64\pi - \frac{8\sqrt{3}}{3} \left[4^{\frac{3}{2}} - 0 \right] - \left[8(0) + 64 \sin^{-1}(1) \right] - \left(4\sqrt{64 - 16} + 64 \sin^{-1} \left(\frac{1}{2} \right) \right)$$

$$= 64\pi - \frac{8\sqrt{3}}{3} (8) - \left[64 \left(\frac{\pi}{2} \right) - 4\sqrt{48} - 64 \frac{\pi}{6} \right]$$



Vikasana - CET 2012



$$= 64\pi - \frac{64\sqrt{3}}{3} - 32\pi + 16\sqrt{3} + \frac{32\pi}{3}$$

$$= \frac{192\pi - 64\sqrt{3} - 96\pi + 48\sqrt{3} + 32\pi}{3}$$

$$= \frac{128\pi - 16\sqrt{3}}{3} = \frac{16}{3} [8\pi - \sqrt{3}] \text{sq. units.}$$

Vikasana - CET 2012



10. The area enclosed between the concyclic circles $x^2+y^2=4$ and $x^2+y^2=9$ is

a) 5π sq. units

b) 4π sq units

c) 9π sq units

d) 36π sq units

Vikasana - CET 2012



Given $x^2 + y^2 = 9 \rightarrow \textcircled{1}$ $x^2 + y^2 = 4 \rightarrow \textcircled{2}$

Let A_1 be the area of circle $\textcircled{1}$ is

$$A_1 = 9\pi \text{ sq. units.}$$

Let A_2 be the area of circle $\textcircled{2}$ is

$$A_2 = 4\pi \text{ sq. units.}$$

Let ' A ' be the area enclosed between the two circles

$$A = A_1 - A_2 = 9\pi - 4\pi \quad \therefore A = 5\pi \text{ sq units}$$

Vikasana - CET 2012



11. Area bounded by the curves

$y = \log x$, $y = \log|x|$, $y = |\log x|$ & $y = |\log|x||$ is

a) 4 sq. units

b) 6 sq. units

c) 10 sq. units

d) none of these

Vikasana - CET 2012



W.K.T. $\log x$ is defined for $x > 0$ and

$\log |x|$ is defined for all $x \in \mathbb{R} - \{0\}$

Also $|\log x| \geq 0$ and $|\log |x|| \geq 0$

Required area is symmetrical in all the four quadrants

$$\text{So the area} = 4 \int_0^1 |\log x| dx$$




$$= -4 \int_0^1 \log x dx \quad [\text{---}]$$

$$= -4 \left[x \log x - x \right]_0^1$$

$$= -4 \left[(1 \log 1 - 1) - (0 - 0) \right]_0^1$$

$$= -4$$





12. The area bounded by the curves $y=x$ & $y=x^3$ is

$$\frac{1}{2} \int_{-1}^1 (x - x^3) dx$$


$$\frac{1}{2} \int_{-1}^1 (x - x^3) dx$$



~~गणित~~ When $x=0, y=0$ $x=\pm 1 \Rightarrow y=\pm 1$
i.e. $x = x^3 \Rightarrow x(x^2 - 1) = 0 \quad \therefore x = 0, x = \pm 1$

\therefore The line $y=x$ intersect the curve $y=x^3$
at three points $(-1,-1), (0,0)$ & $(1,1)$ Hence
it is symmetric in opposite quadrant.

~~गणित~~



Vikasana - CET 2012



$$= \int_0^1 (x - x^3) dx + \int_{-1}^0 (x^3 - x) dx$$

$$= \left[\frac{x^2}{2} - \frac{x^4}{4} \right]_0^1 + \left[\frac{x^4}{4} - \frac{x^2}{2} \right]_{-1}^0$$

$$= \frac{1}{4} + \frac{1}{4} = \frac{1}{2} \text{ sq units}$$

Vikasana - CET 2012



13. The area bounded by the curves $|x| + |y| \geq 1$ and $x^2 + y^2 \leq 1$ is

a) $2\sqrt{2}$

b) $\pi - 2$

c) $2\sqrt{2} - \pi$

d) $\pi - 2\sqrt{2}$



14) The area of region bounded by $x^2=16y$ & $x=0$ and $y=1, y=4$ and y -axis in the 1st quadrant is

$$\frac{64}{3} \text{ sq. units}$$




$$\frac{56}{3} \text{ sq. units}$$

$$\frac{16}{3} \text{ sq. units}$$

$$\frac{4}{3} \text{ sq. units}$$



$\int_1^4 \frac{1}{x} dx$



Vikasana - CET 2012



15) The area of the region bounded by $y=x^2-5x+4$ and x -axis is

a) $\frac{3}{2}$

b) $\frac{5}{2}$

c) $\frac{7}{2}$

d) $\frac{9}{2}$



Since the curve $y = x^2 - 5x + 4$
crosses x -axis $y=0$

~~$x^2 - 5x + 4 = 0$~~ $x^2 - 4x - 1x + 4 = 0$

$(x-4)(x-1) = 0 \quad \therefore x = 1, 4$




~~Repeat~~ $\int_1^4 y \, dx :$

$\int_1^4 (x^2 - 5x + 4) \, dx$

$\left[\frac{x^3}{3} - \frac{5x^2}{2} + 4x \right]_1^4$

Vikasana - CET 2012





$$= \left(\frac{64}{3} - \frac{80}{2} + 16 \right) - \left(\frac{1}{3} - \frac{5}{2} + 4 \right)$$

$$= \left(\frac{64}{3} - 24 \right) - \left(\frac{2 - 15 + 24}{6} \right)$$

$$= \left(\frac{64 - 72}{3} \right) - \left(\frac{11}{6} \right)$$

$$= \left| \frac{-16 - 11}{6} \right| = \frac{27}{6} = \frac{9}{2}$$

Vikasana - CET 2012



16. The area enclosed by the parabola $y^2=16x$ and its latus rectum

$$\frac{1}{3} \times 108$$

$$\frac{1}{3} \times 16$$

$$\frac{1}{3} \times 128$$

$$\frac{4}{3}$$




भारतीय

$$2 \int_0^4 y dx$$

$$= 2 \int_0^4 4\sqrt{x} dx = 8 \left[\frac{x^{\frac{3}{2}}}{\frac{3}{2}} \right]_0^4$$

$$= \frac{16}{3} \left[4^{\frac{3}{2}} - 0 \right] = \frac{16}{3} (8)$$



128
3

Vikasana - CET 2012



17. The area of smaller segment cut off from the circle $x^2+y^2=9$ by $x=1$ is

~~$\frac{1}{2} [9\sec^{-1} \frac{1}{3} - \sqrt{8}]$~~ b) $(9\sec^{-1} 3 - \sqrt{8})$ sq units

c) $(\sqrt{8} - 9\sec^{-1} 3)$ sq units ~~$\frac{1}{2} [9\sec^{-1} \frac{1}{3} - \sqrt{8}]$~~




$$R_{\text{area}} = 2 \int_1^3 y \, dx = 2 \int_1^3 \sqrt{9-x^2} \, dx$$

$$= 2 \left[\frac{x\sqrt{9-x^2}}{2} + \frac{9}{2} \sin^{-1} \left(\frac{x}{3} \right) \right]_1^3$$

$$= \left[\sqrt{9-9} + \sqrt{9-1} \right] - \left[\sqrt{9-1} + \sqrt{9-9} \right]$$

$$= 9 \left(\frac{\pi}{2} \right) - \sqrt{8} - 9 \sin^{-1} \frac{1}{3} = 9 \left(\frac{\pi}{2} - \sin^{-1} \left(\frac{1}{3} \right) - \sqrt{8} \right)$$

$$= 9 \cos^{-1} \left(\frac{1}{3} \right) - \sqrt{8} = 9 \operatorname{Sec}^{-1} 3 - \sqrt{8}$$



Vikasana - CET 2012



18. The ratio of which the area bounded by the curves $y^2=12x$ and $x^2=12y$ is divided by the line $x=3$ is

a) 154

b) 134

c) 123

d) None




$$\text{Let } A = \int_0^3 \sqrt{12x} \, dx - \int_0^3 \frac{x^2}{2} \, dx$$

$$= \left[2\sqrt{3} \frac{x^{\frac{3}{2}}}{\frac{3}{2}} - \frac{x^3}{36} \right]_0^3$$

$$= \frac{4\sqrt{3}}{3} (3\sqrt{3}) - \frac{1}{36} (27) = 12\frac{3}{4}$$

45
45
45
45

Vikasana - CET 2012





$$\text{Let } A_2 = \int_3^{12} \sqrt{12x} \, dx - \int_3^{12} \frac{x^2}{12} \, dx$$

$$= \frac{4\sqrt{3}}{3} \left[12^{\frac{3}{2}} - 3^{\frac{3}{2}} \right] - \frac{1}{36} (12^3 - 3^3)$$

$$= \frac{4\sqrt{3}}{3} [24\sqrt{3} - 3\sqrt{3}] - \frac{1}{36} (1728 - 27)$$

$$= \frac{4\sqrt{3}}{3} [21\sqrt{3}] - \frac{1}{36} (1701) = 84 - \frac{1701}{36} = 84 - \frac{189}{4} = \frac{336}{4} - \frac{189}{4} = \frac{147}{4}$$



Vikasana - CET 2012



19. The area bounded by $y=ax^2$ and $x=ay^2$ ($a>0$) is 1 then 'a' is

a) 1

b) $\frac{1}{\sqrt{3}}$

c) $\frac{1}{3}$

d) $\frac{-1}{\sqrt{3}}$




Solve the given equations, we get $(0,0)$ & $\left(\frac{1}{a}, \frac{1}{a}\right)$

~~By A=~~ Area of $OCBDO$ – Area of $OABDO$

$$1 = \int_0^{\frac{1}{a}} \left[\sqrt{x} - \sqrt{a} \right] dx$$

$$\Rightarrow 1 = \left[\frac{1}{\sqrt{a}} \frac{x^{\frac{3}{2}}}{\frac{3}{2}} - a \cdot \frac{x^3}{3} \right]_0^{\frac{1}{a}}$$



Vikasana - CET 2012



$$I = \begin{bmatrix} 2 & 1 & 0 \\ \sqrt{3} & \sqrt{3} & \sqrt{3} \end{bmatrix} \begin{bmatrix} a & 1 \\ 3 & a \end{bmatrix} \epsilon$$

$$I = \begin{bmatrix} 2 & 1 \\ \sqrt{3} & \sqrt{3} \end{bmatrix}$$

$$I = \begin{bmatrix} 1 & 1 \\ \sqrt{3} & \sqrt{3} \end{bmatrix} \begin{bmatrix} 1 & 1 \\ \sqrt{3} & \sqrt{3} \end{bmatrix} \epsilon$$

Vikasana - CET 2012



20. The area of the region
 $\{ (x, y): x^2 + y^2 \leq 1 \leq x + y \}$

$$\frac{\pi}{5} \text{ Sq.} \quad \frac{\pi}{4} \text{ Sq.}$$

$$\frac{\pi}{3} \text{ Sq.} \quad \frac{\pi}{4} \left(\frac{1}{2} \right) \text{ Sq.}$$



Given equation of the circle and the line are $x^2 + y^2 = 1$ and $x + y = 1$

Solving these equations we get


$$x=0, x=1$$

$A(1,0)$ and $B(0,1)$

Required Area =

Area of OAB - Area of triangle OAB

$$= \left(\frac{\pi}{4} - \frac{1}{2} \right) \text{Sq. units}$$



Vikasana - CET 2012



21. Area of included between the curves $y=x^2-3x+2$ and $y=-x^2+3x-2$ is

~~a) $\frac{1}{6}$ sq unit~~ ~~b) $\frac{1}{2}$ sq unit~~

~~c) 1 sq unit~~ d) $\frac{1}{3}$ sq unit



~~$\int (x^3 - 3x^2 + 2x) dx$~~

*Bojortti
arsame*

$$\begin{aligned} &= 2 \left(\frac{x^3}{3} - \frac{3x^2}{2} + 2x \right)_1^2 \\ &= 2 \left(\frac{8}{3} - 6 + 4 \right) - \left(\frac{1}{3} - \frac{3}{2} + 2 \right) \\ &= 2 \left(\frac{8}{3} - 2 \right) - \left(\frac{2 - 9 + 12}{6} \right) \\ &= 2 \left[\frac{2}{3} - \frac{5}{6} \right] = 2 \left[\frac{1}{6} \right] = \frac{1}{3} \text{ squnits} \end{aligned}$$

Vikasana - CET 2012



22. The area bounded by the curve $y=e^{|x|}$, x -axis and the lines $x=-1$ and $x=1$ is

~~2e - 2~~ 2e - 2

~~2e - 2~~ 2e - 2




$$R_{\text{area}} = \int_{-1}^{+1} e^{|x|} dx$$

$$= 2 \int_0^1 e^x dx$$

$$= 2 \left[e^x \right]_0^1$$

$$= 2 \left[e^1 - e^0 \right]$$

$$= 2 \left[e - 1 \right] \text{sq units}$$



Vikasana - CET 2012



23. The area bounded by the curve $x^2 = y + 4$ and the lines $y = 0$ and $y = 5$ is

$$\frac{16}{3}$$

$$\frac{76}{3}$$

$$\frac{20}{3}$$

$$\frac{1}{3}$$



Reqd A = Area of ABCDA = 2(Area of ABMOA)


$$= 2 \int_0^5 x \, dy$$

$$= 2 \int_0^5 \sqrt{y+4} \, dy$$

$$= 2 \left[\frac{(y+4)^{3/2}}{3/2} \right]_0^5$$

$$= \frac{4}{3} [27 - 8] = \frac{4}{3} [9^{3/2} - 4^{3/2}] = \frac{4}{3} [19]$$

$$A = \frac{76}{3} \text{ squnits}$$



Vikasana - CET 2012



24. The area region bounded by $x = a \cos \theta$

and $y = a \sin \theta$ or $x = a \left[\frac{1 - t^2}{1 + t^2} \right]$ &

$y = a \left[\frac{2t}{1 + t^2} \right]$ is

a) $2\pi a^2$

b) πa^2

c) 2π

d) π

Vikasana - CET 2012



25. The area of the region bounded by $x = a \cos \theta$ and $y = b \sin \theta$, i.e.

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \quad (a > b) \text{ is}$$

a) $2\pi a$

b) πa

c) $4\pi a$

d) ab

Vikasana - CET 2012